

My healthcare professional recommends...

Children's Advil® Relief you can trust

Children's Advil®

Lasts Up To

8
Hours

Find the right dose on the chart below. If possible, **use weight to dose** otherwise use age.



Infants' Advil® Concentrated Drops for ages 6-23 months

ACTIVE INGREDIENT
Ibuprofen 50 mg (NSAID)[†] in each 1.25 mL

Fever reducer; relieves minor aches and pains due to the common cold, flu, headaches, and toothaches



Children's Advil® Suspension for ages 2-11 years

ACTIVE INGREDIENT
Ibuprofen 100 mg (NSAID)[†] in each 5 mL

Fever reducer; relieves minor aches and pains due to the common cold, flu, sore throat headaches, and toothaches

Available in 4 flavors!
• Dye-Free White Grape
• Fruit
• Grape
• Blue Raspberry



Junior Strength Advil® Chewables for ages 6-11 years

ACTIVE INGREDIENT
Ibuprofen 100 mg (NSAID)[†] in each tablet

Fever reducer; relieves minor aches and pains due to the common cold, flu, sore throat headaches, and toothaches



Junior Strength Advil® Tablets for ages 6-11 years

ACTIVE INGREDIENT
Ibuprofen 100 mg (NSAID)[†] in each tablet

Fever reducer; relieves minor aches and pains due to the common cold, flu, sore throat headaches, and toothaches

Weight		Age	GIVE YOUR CHILD THE DOSE INDICATED BELOW EVERY 6-8 HOURS, NOT TO EXCEED 4 DOSES/24 HOURS							
PATIENT DOSING	12-17 pounds	under 6 months	Ask a doctor		Ask a doctor about using Infants' Advil®					
		6-11 months	1.25 mL [‡]							
	18-23 pounds	12-23 months	1.875 mL [‡]				Ask a doctor about using Infants' or Children's Advil®			
	24-35 pounds	2-3 years	Ask a doctor about using Children's or Junior Strength Advil®						1 tsp [‡]	
	36-47 pounds	4-5 years			1½ tsp [‡]					
	48-59 pounds	6-8 years			2 tsp [‡]				2 tablets	2 tablets
	60-71 pounds	9-10 years			2½ tsp [‡]				2½ tablets	2 tablets
	72-95 pounds	11 years	3 tsp [‡]		3 tablets	3 tablets				

For 12+ years, use adult strength Advil®

Use the fever diary below to record your child's fevers and dosing schedule.

Fever:	_____°F or _____°C	Time:	_____	Dose:	_____
Fever:	_____°F or _____°C	Time:	_____	Dose:	_____
Fever:	_____°F or _____°C	Time:	_____	Dose:	_____
Fever:	_____°F or _____°C	Time:	_____	Dose:	_____

Please refer to Drug Facts on the product label or visit Advil.com for complete warnings.

Important Instructions for Proper Use

- Read and keep the cartons for complete warnings and information on Advil® products
- If you are taking other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), then consult your doctor before taking any Advil® products
- Adult strength medicines should not be administered to children
- Measure dosing only with the device that comes with the product

Pfizer Consumer Healthcare

[†]Nonsteroidal anti-inflammatory drug.
[‡]Measure dosing only with the device provided.
Shake liquid products well before dosing.

Visit ChildrensAdvil.com for Drug Facts, product updates and additional information.