

Dosing for Infants and Children

From Your Healthcare Professional

DOSE: Every 6-8 hours as needed. DO NOT GIVE MORE THAN 4 DOSES IN 24 HOURS.

If possible, use weight to dose; otherwise use age.

mL = milliliter
tsp = teaspoon



Infants' MOTRIN® Concentrated Drops

Active Ingredient:
Ibuprofen 50 mg (NSAID)[†]
(in each 1.25 mL)

[†]Nonsteroidal anti-inflammatory drug
Use only as directed.

Available in:
 Dye-Free Berry



Children's MOTRIN® Oral Suspension

Active Ingredient:
Ibuprofen 100 mg (NSAID)[†]
(in each 5 mL or 1 tsp)

[†]Nonsteroidal anti-inflammatory drug
Use only as directed.

Available in:
 Original Berry
 Dye-Free Berry

WEIGHT	AGE	Infants' MOTRIN® Concentrated Drops	Children's MOTRIN® Oral Suspension
6-11 lbs	0-5 mos	Do not use	—
12-17 lbs	6-11 mos	1.25 mL 	—
18-23 lbs	12-23 mos	1.875 mL 	—
24-35 lbs	2-3 yrs	—	5 mL (1 tsp)
36-47 lbs	4-5 yrs	—	7.5 mL (1½ tsp)
48-59 lbs	6-8 yrs	—	10 mL (2 tsp)
60-71 lbs	9-10 yrs	—	12.5 mL (2½ tsp)
72-95 lbs	11 yrs	—	15 mL (3 tsp)

IMPORTANT INSTRUCTIONS for Proper Use

Today's Date: _____

This dosing recommendation from your doctor will expire in 14 DAYS.


- Read and follow the label on all MOTRIN® products.
- Take every 6-8 hours as needed. Do NOT exceed more than 4 doses in 24 hours.
- Do NOT administer longer than 10 days, unless directed by a doctor.
- Ask a doctor or pharmacist before use if the child is taking any other drug containing an NSAID[†] (prescription or non-prescription).
— MOTRIN® contains ibuprofen
- Use only the dosing device that comes with a specific product.

[†]Nonsteroidal anti-inflammatory drug



Kids' Wellness Tracker

This free, all-in-one tool from the makers of Children's TYLENOL® and Children's MOTRIN® makes it easy to track your child's height, weight, vaccines, symptoms, and medicines, plus calculate dosing.



Download now at the App StoreSM

App Store is a service mark of Apple Inc. Standard download rates may apply